

CAPSULES, ORAL SUSPENSION  
and SUPPOSITORIES

**INDOCIN®**  
**(INDOMETHACIN)**

If you take Indocin to relieve gout pain,  
please read the [Indocin For Gout Guidelines](#)  
and other indomethacin pages at  
[GoutPal.com](#)

#### **Cardiovascular Risk**

- NSAIDs may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at a greater risk. (See **WARNINGS**.)
- INDOCIN is contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery. (See **WARNINGS**.)

#### **Gastrointestinal Risk**

- NSAIDs cause an increased risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious gastrointestinal events. (See **WARNINGS**.)

#### **DESCRIPTION**

INDOCIN\* is supplied in three dosage forms. Capsules INDOCIN for oral administration contain either 25 mg or 50 mg of indomethacin and the following inactive ingredients: colloidal silicon dioxide, FD&C Blue 1, FD&C Red 3, gelatin, lactose, lecithin, magnesium stearate, and titanium dioxide. Suspension INDOCIN for oral use contains 25 mg of indomethacin per 5 mL, alcohol 1%, and sorbic acid 0.1% added as a preservative and the following inactive ingredients: antifoam AF emulsion, flavors, purified water, sodium hydroxide or hydrochloric acid to adjust pH, sorbitol solution, and tragacanth. Suppositories INDOCIN for rectal use contain 50 mg of indomethacin and the following inactive ingredients: butylated hydroxyanisole, butylated hydroxytoluene, edetic acid, glycerin, polyethylene glycol 3350, polyethylene glycol 8000 and sodium chloride. Indomethacin is a non-steroidal anti-inflammatory indole derivative designated chemically as 1-(4-chlorobenzoyl)-5-methoxy-2-methyl-1*H*-indole-3-acetic acid. Indomethacin is practically insoluble in water and sparingly soluble in alcohol. It has a pKa of 4.5 and is stable in neutral or slightly acidic media and decomposes in strong alkali. The suspension has a pH of 4.0-5.0. The structural formula is:

\* Registered trademark of MERCK & CO., Inc.  
COPYRIGHT © 1988, 2005 MERCK & CO., Inc.  
All rights reserved

## CLINICAL PHARMACOLOGY

INDOCIN is a non-steroidal anti-inflammatory drug (NSAID) that exhibits antipyretic and analgesic properties. Its mode of action, like that of other anti-inflammatory drugs, is not known. However, its therapeutic action is not due to pituitary-adrenal stimulation.

INDOCIN is a potent inhibitor of prostaglandin synthesis *in vitro*. Concentrations are reached during therapy which have been demonstrated to have an effect *in vivo* as well. Prostaglandins sensitize afferent nerves and potentiate the action of bradykinin in inducing pain in animal models. Moreover, prostaglandins are known to be among the mediators of inflammation. Since indomethacin is an inhibitor of prostaglandin synthesis, its mode of action may be due to a decrease of prostaglandins in peripheral tissues.

INDOCIN has been shown to be an effective anti-inflammatory agent, appropriate for long-term use in rheumatoid arthritis, ankylosing spondylitis, and osteoarthritis.

INDOCIN affords relief of symptoms; it does not alter the progressive course of the underlying disease.

INDOCIN suppresses inflammation in rheumatoid arthritis as demonstrated by relief of pain, and reduction of fever, swelling and tenderness. Improvement in patients treated with INDOCIN for rheumatoid arthritis has been demonstrated by a reduction in joint swelling, average number of joints involved, and morning stiffness; by increased mobility as demonstrated by a decrease in walking time; and by improved functional capability as demonstrated by an increase in grip strength. INDOCIN may enable the reduction of steroid dosage in patients receiving steroids for the more severe forms of rheumatoid arthritis. In such instances the steroid dosage should be reduced slowly and the patients followed very closely for any possible adverse effects.

Indomethacin has been reported to diminish basal and CO<sub>2</sub> stimulated cerebral blood flow in healthy volunteers following acute oral and intravenous administration. In one study after one week of treatment with orally administered indomethacin, this effect on basal cerebral blood flow had disappeared. The clinical significance of this effect has not been established.

Capsules INDOCIN have been found effective in relieving the pain, reducing the fever, swelling, redness, and tenderness of acute gouty arthritis — see **INDICATIONS AND USAGE**.

Following single oral doses of Capsules INDOCIN 25 mg or 50 mg, indomethacin is readily absorbed, attaining peak plasma concentrations of about 1 and 2 mcg/mL, respectively, at about 2 hours. Orally administered Capsules INDOCIN are virtually 100% bioavailable, with 90% of the dose absorbed within 4 hours. A single 50 mg dose of Oral Suspension INDOCIN was found to be bioequivalent to a 50 mg INDOCIN capsule when each was administered with food.

Indomethacin is eliminated via renal excretion, metabolism, and biliary excretion. Indomethacin undergoes appreciable enterohepatic circulation. The mean half-life of indomethacin is estimated to be about 4.5 hours. With a typical therapeutic regimen of 25 or 50 mg t.i.d., the steady-state plasma concentrations of indomethacin are an average 1.4 times those following the first dose.

The rate of absorption is more rapid from the rectal suppository than from Capsules INDOCIN. Ordinarily, therefore, the total amount absorbed from the suppository would be expected to be at least equivalent to the capsule. In controlled clinical trials, however, the amount of indomethacin absorbed was found to be somewhat less (80-90%) than that absorbed from Capsules INDOCIN. This is probably because some subjects did not retain the material from the suppository for the one hour necessary to assure complete absorption. Since the suppository dissolves rather quickly rather than melting slowly, it is seldom recovered in recognizable form if the patient retains the suppository for more than a few minutes.

Indomethacin exists in the plasma as the parent drug and its desmethyl, desbenzoyl, and desmethyl-desbenzoyl metabolites, all in the unconjugated form. About 60 percent of an oral dosage is recovered in urine as drug and metabolites (26 percent as indomethacin and its glucuronide), and 33 percent is recovered in feces (1.5 percent as indomethacin).

About 99% of indomethacin is bound to protein in plasma over the expected range of therapeutic plasma concentrations. Indomethacin has been found to cross the blood-brain barrier and the placenta.

## INDICATIONS AND USAGE

Carefully consider the potential benefits and risks of INDOCIN and other treatment options before deciding to use INDOCIN. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

Indomethacin has been found effective in active stages of the following:

1. Moderate to severe rheumatoid arthritis including acute flares of chronic disease.
2. Moderate to severe ankylosing spondylitis.
3. Moderate to severe osteoarthritis.
4. Acute painful shoulder (bursitis and/or tendinitis).
5. Acute gouty arthritis.

## CONTRAINDICATIONS

INDOCIN is contraindicated in patients with known hypersensitivity to indomethacin or the excipients (see **DESCRIPTION**).

INDOCIN should not be given to patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Severe, rarely fatal, anaphylactic/anaphylactoid reactions to NSAIDs have been reported in such patients (see **WARNINGS – Anaphylactic/Anaphylactoid Reactions**, and **PRECAUTIONS – Preexisting Asthma**).

INDOCIN is contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery (see **WARNINGS**).

Suppositories INDOCIN are contraindicated in patients with a history of proctitis or recent rectal bleeding.

## WARNINGS

### CARDIOVASCULAR EFFECTS

#### Cardiovascular Thrombotic Events

Clinical trials of several COX-2 selective and nonselective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, myocardial infarction, and stroke, which can be fatal. All NSAIDs, both COX-2 selective and nonselective, may have a similar risk. Patients with known CV disease or risk factors for CV disease may be at greater risk. To minimize the potential risk for an adverse CV event in patients treated with an NSAID, the lowest effective dose should be used for the shortest duration possible. Physicians and patients should remain alert for the development of such events, even in the absence of previous CV symptoms. Patients should be informed about the signs and/or symptoms of serious CV events and the steps to take if they occur.

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID does increase the risk of serious GI events (see **GI WARNINGS**).

Two large, controlled, clinical trials of a COX-2 selective NSAID for the treatment of pain in the first 10-14 days following CABG surgery found an increased incidence of myocardial infarction and stroke (see **CONTRAINDICATIONS**).

#### Hypertension

NSAIDs, including INDOCIN, can lead to onset of new hypertension or worsening of pre-existing hypertension, either of which may contribute to the increased incidence of CV events. Patients taking thiazides or loop diuretics may have impaired response to these therapies when taking NSAIDs. NSAIDs, including INDOCIN, should be used with caution in patients with hypertension. Blood pressure (BP) should be monitored closely during the initiation of NSAID treatment and throughout the course of therapy.

**Congestive Heart Failure and Edema**

Fluid retention and edema have been observed in some patients taking NSAIDs. INDOCIN should be used with caution in patients with fluid retention or heart failure.

In a study of patients with severe heart failure and hyponatremia, INDOCIN was associated with significant deterioration of circulatory hemodynamics, presumably due to inhibition of prostaglandin dependent compensatory mechanisms.

**Gastrointestinal Effects – Risk of Ulceration, Bleeding, and Perforation**

NSAIDs, including INDOCIN, can cause serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the esophagus, stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients, who develop a serious upper GI adverse event on NSAID therapy is symptomatic. Upper GI ulcers, gross bleeding, or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3-6 months, and in about 2-4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious GI event at some time during the course of therapy. However, even short-term therapy is not without risk.

Rarely, in patients taking INDOCIN, intestinal ulceration has been associated with stenosis and obstruction. Gastrointestinal bleeding without obvious ulcer formation and perforation of pre-existing sigmoid lesions (diverticulum, carcinoma, etc.) have occurred. Increased abdominal pain in ulcerative colitis patients or the development of ulcerative colitis and regional ileitis have been reported to occur rarely.

NSAIDs should be prescribed with extreme caution in those with prior history of ulcer disease or gastrointestinal bleeding. Patients with a *prior history of peptic ulcer disease and/or gastrointestinal bleeding* who use NSAIDs have a greater than 10-fold increased risk for developing a GI bleed compared to patients with neither of these risk factors. Other factors that increase the risk for GI bleeding in patients treated with NSAIDs include concomitant use of oral corticosteroids or anticoagulants, longer duration of NSAID therapy, smoking, use of alcohol, older age, and poor general health status. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and therefore, special care should be taken in treating this population.

To minimize the potential risk for an adverse GI event in patients treated with an NSAID, the lowest effective dose should be used for the shortest possible duration. Patients and physicians should remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected. This should include discontinuation of the NSAID until a serious GI adverse event is ruled out. For high risk patients, alternate therapies that do not involve NSAIDs should be considered.

**Renal Effects**

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of a nonsteroidal anti-inflammatory drug may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate over renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors, patients with volume depletion, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state.

Increases in serum potassium concentration, including hyperkalemia, have been reported with use of INDOCIN, even in some patients without renal impairment. In patients with normal renal function, these effects have been attributed to a hyporeninemic-hypoaldosteronism state (see **PRECAUTIONS, Drug Interactions**).

**Advanced Renal Disease**

No information is available from controlled clinical studies regarding the use of INDOCIN in patients with advanced renal disease. Therefore, treatment with INDOCIN is not recommended in these patients with advanced renal disease. If INDOCIN therapy must be initiated, close monitoring of the patient's renal function is advisable.

**Anaphylactic/Anaphylactoid Reactions**

As with other NSAIDs, anaphylactic/anaphylactoid reactions may occur in patients without known prior exposure to INDOCIN. INDOCIN should not be given to patients with the aspirin triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking aspirin or other NSAIDs (see

**CONTRAINDICATIONS** and **PRECAUTIONS – Preexisting Asthma**). Emergency help should be sought in cases where an anaphylactic/anaphylactoid reaction occurs.

**Skin Reactions**

NSAIDs, including INDOCIN, can cause serious skin adverse events such as exfoliative dermatitis, Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. These serious events may occur without warning. Patients should be informed about the signs and symptoms of serious skin manifestations and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

**Pregnancy**

In late pregnancy, as with other NSAIDs, INDOCIN should be avoided because it may cause premature closure of the ductus arteriosus.

**Ocular Effects**

Corneal deposits and retinal disturbances, including those of the macula, have been observed in some patients who had received prolonged therapy with INDOCIN. The prescribing physician should be alert to the possible association between the changes noted and INDOCIN. It is advisable to discontinue therapy if such changes are observed. Blurred vision may be a significant symptom and warrants a thorough ophthalmological examination. Since these changes may be asymptomatic, ophthalmologic examination at periodic intervals is desirable in patients where therapy is prolonged.

**Central Nervous System Effects**

INDOCIN may aggravate depression or other psychiatric disturbances, epilepsy, and parkinsonism, and should be used with considerable caution in patients with these conditions. If severe CNS adverse reactions develop, INDOCIN should be discontinued.

INDOCIN may cause drowsiness; therefore, patients should be cautioned about engaging in activities requiring mental alertness and motor coordination, such as driving a car. INDOCIN may also cause headache. Headache which persists despite dosage reduction requires cessation of therapy with INDOCIN.

**PRECAUTIONS****General**

INDOCIN cannot be expected to substitute for corticosteroids or to treat corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to disease exacerbation. Patients on prolonged corticosteroid therapy should have their therapy tapered slowly if a decision is made to discontinue corticosteroids.

The pharmacological activity of INDOCIN in reducing fever and inflammation may diminish the utility of these diagnostic signs in detecting complications of presumed noninfectious, painful conditions.

**Hepatic Effects**

Borderline elevations of one or more liver tests may occur in up to 15% of patients taking NSAIDs including INDOCIN. These laboratory abnormalities may progress, may remain unchanged, or may be transient with continuing therapy. Notable elevations of ALT or AST (approximately three or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials with NSAIDs. In addition, rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis and hepatic failure, some of them with fatal outcomes have been reported.

A patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of a more severe hepatic reaction while on therapy with INDOCIN. If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash, etc.), INDOCIN should be discontinued.

**Hematological Effects**

Anemia is sometimes seen in patients receiving NSAIDs, including INDOCIN. This may be due to fluid retention, occult or gross GI blood loss, or an incompletely described effect upon erythropoiesis. Patients on long-term treatment with NSAIDs, including INDOCIN, should have their hemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia.

NSAIDs inhibit platelet aggregation and have been shown to prolong bleeding time in some patients. Unlike aspirin, their effect on platelet function is quantitatively less, of shorter duration, and reversible. Patients receiving INDOCIN who may be adversely affected by alterations in platelet function, such as those with coagulation disorders or patients receiving anticoagulants, should be carefully monitored.

**Preexisting Asthma**

Patients with asthma may have aspirin-sensitive asthma. The use of aspirin in patients with aspirin-sensitive asthma has been associated with severe bronchospasm which can be fatal. Since cross reactivity, including bronchospasm, between aspirin and other nonsteroidal anti-inflammatory drugs has been reported in such aspirin-sensitive patients, INDOCIN should not be administered to patients with this form of aspirin sensitivity and should be used with caution in patients with preexisting asthma.

**Information for Patients**

**Patients should be informed of the following information before initiating therapy with an NSAID and periodically during the course of ongoing therapy. Patients should also be encouraged to read the NSAID Medication Guide that accompanies each prescription dispensed.**

1. INDOCIN, like other NSAIDs, may cause serious CV side effects, such as MI or stroke, which may result in hospitalization and even death. Although serious CV events can occur without warning symptoms, patients should be alert for the signs and symptoms of chest pain, shortness of breath, weakness, slurring of speech, and should ask for medical advice when observing any indicative sign or symptoms. Patients should be apprised of the importance of this follow-up (see **WARNINGS, CARDIOVASCULAR EFFECTS**).
2. INDOCIN, like other NSAIDs, can cause GI discomfort and, rarely, serious GI side effects, such as ulcers and bleeding, which may result in hospitalization and even death. Although serious GI tract ulcerations and bleeding can occur without warning symptoms, patients should be alert for the signs and symptoms of ulcerations and bleeding, and should ask for medical advice when observing any indicative sign or symptoms including epigastric pain, dyspepsia, melena, and hematemesis. Patients should be apprised of the importance of this follow-up (see **WARNINGS, Gastrointestinal Effects - Risk of Ulceration, Bleeding, and Perforation**).
3. INDOCIN, like other NSAIDs, can cause serious skin side effects such as exfoliative dermatitis, SJS, and TEN, which may result in hospitalizations and even death. Although serious skin reactions may occur without warning, patients should be alert for the signs and symptoms of skin rash and blisters, fever, or other signs of hypersensitivity such as itching, and should ask for medical advice when observing any indicative signs or symptoms. Patients should be advised to stop the drug immediately if they develop any type of rash and contact their physicians as soon as possible.
4. Patients should promptly report signs or symptoms of unexplained weight gain or edema to their physicians.
5. Patients should be informed of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, pruritus, jaundice, right upper quadrant tenderness, and “flu-like” symptoms). If these occur, patients should be instructed to stop therapy and seek immediate medical therapy.
6. Patients should be informed of the signs of an anaphylactic/anaphylactoid reaction (e.g. difficulty breathing, swelling of the face or throat). If these occur, patients should be instructed to seek immediate emergency help (see **WARNINGS**).

7. In late pregnancy, as with other NSAIDs, INDOCIN should be avoided because it may cause premature closure of the ductus arteriosus.

**Laboratory Tests**

Because serious GI tract ulcerations and bleeding can occur without warning symptoms, physicians should monitor for signs or symptoms of GI bleeding. Patients on long-term treatment with NSAIDs should have their CBC and a chemistry profile checked periodically. If clinical signs and symptoms consistent with liver or renal disease develop, systemic manifestations occur (e.g., eosinophilia, rash, etc.) or if abnormal liver tests persist or worsen, INDOCIN should be discontinued.

**Drug Interactions****ACE-Inhibitors and Angiotensin II Antagonists**

Reports suggest that NSAIDs may diminish the antihypertensive effect of ACE-inhibitors and angiotensin II antagonists. INDOCIN can reduce the antihypertensive effects of captopril and losartan. These interactions should be given consideration in patients taking NSAIDs concomitantly with ACE-inhibitors or angiotensin II antagonists. In some patients with compromised renal function, the co-administration of an NSAID and an ACE-inhibitor or an angiotensin II antagonist may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible.

**Aspirin**

When INDOCIN is administered with aspirin, its protein binding is reduced, although the clearance of free INDOCIN is not altered. The clinical significance of this interaction is not known.

The use of INDOCIN in conjunction with aspirin or other salicylates is not recommended. Controlled clinical studies have shown that the combined use of INDOCIN and aspirin does not produce any greater therapeutic effect than the use of INDOCIN alone. In a clinical study of the combined use of INDOCIN and aspirin, the incidence of gastrointestinal side effects was significantly increased with combined therapy.

In a study in normal volunteers, it was found that chronic concurrent administration of 3.6 g of aspirin per day decreases indomethacin blood levels approximately 20%.

**Beta-adrenoceptor blocking agents**

Blunting of the antihypertensive effect of beta-adrenoceptor blocking agents by non-steroidal anti-inflammatory drugs including INDOCIN has been reported. Therefore, when using these blocking agents to treat hypertension, patients should be observed carefully in order to confirm that the desired therapeutic effect has been obtained.

**Cyclosporine**

Administration of non-steroidal anti-inflammatory drugs concomitantly with cyclosporine has been associated with an increase in cyclosporine-induced toxicity, possibly due to decreased synthesis of renal prostacyclin. NSAIDs should be used with caution in patients taking cyclosporine, and renal function should be carefully monitored.

**Diflunisal**

In normal volunteers receiving indomethacin, the administration of diflunisal decreased the renal clearance and significantly increased the plasma levels of indomethacin. In some patients, combined use of INDOCIN and diflunisal has been associated with fatal gastrointestinal hemorrhage. Therefore, diflunisal and INDOCIN should not be used concomitantly.

**Digoxin**

INDOCIN given concomitantly with digoxin has been reported to increase the serum concentration and prolong the half-life of digoxin. Therefore, when INDOCIN and digoxin are used concomitantly, serum digoxin levels should be closely monitored.

**Diuretics**

In some patients, the administration of INDOCIN can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing, and thiazide diuretics. This response has been attributed to inhibition of renal prostaglandin synthesis.

INDOCIN reduces basal plasma renin activity (PRA), as well as those elevations of PRA induced by furosemide administration, or salt or volume depletion. These facts should be considered when evaluating plasma renin activity in hypertensive patients.

It has been reported that the addition of triamterene to a maintenance schedule of INDOCIN resulted in reversible acute renal failure in two of four healthy volunteers. INDOCIN and triamterene should not be administered together.

INDOCIN and potassium-sparing diuretics each may be associated with increased serum potassium levels. The potential effects of INDOCIN and potassium-sparing diuretics on potassium kinetics and renal function should be considered when these agents are administered concurrently.

Most of the above effects concerning diuretics have been attributed, at least in part, to mechanisms involving inhibition of prostaglandin synthesis by INDOCIN.

During concomitant therapy with NSAIDs, the patient should be observed closely for signs of renal failure (see **WARNINGS, Renal Effects**), as well as to assure diuretic efficacy.

### **Lithium**

Capsules INDOCIN 50 mg t.i.d. produced a clinically relevant elevation of plasma lithium and reduction in renal lithium clearance in psychiatric patients and normal subjects with steady state plasma lithium concentrations. This effect has been attributed to inhibition of prostaglandin synthesis. As a consequence, when NSAIDs and lithium are given concomitantly, the patient should be carefully observed for signs of lithium toxicity. (Read circulars for lithium preparations before use of such concomitant therapy.) In addition, the frequency of monitoring serum lithium concentration should be increased at the outset of such combination drug treatment.

### **Methotrexate**

NSAIDs have been reported to competitively inhibit methotrexate accumulation in rabbit kidney slices. This may indicate that they could enhance the toxicity of methotrexate. Caution should be used when NSAIDs are administered concomitantly with methotrexate.

### **NSAIDs**

The concomitant use of INDOCIN with other NSAIDs is not recommended due to the increased possibility of gastrointestinal toxicity, with little or no increase in efficacy.

### **Oral anticoagulants**

Clinical studies have shown that INDOCIN does not influence the hypoprothrombinemia produced by anticoagulants. However, when any additional drug, including INDOCIN, is added to the treatment of patients on anticoagulant therapy, the patients should be observed for alterations of the prothrombin time. In post-marketing experience, bleeding has been reported in patients on concomitant treatment with anticoagulants and INDOCIN. Caution should be exercised when INDOCIN and anticoagulants are administered concomitantly. The effects of warfarin and NSAIDs on GI bleeding are synergistic, such that users of both drugs together have a risk of serious GI bleeding higher than users of either drug alone.

### **Probenecid**

When INDOCIN is given to patients receiving probenecid, the plasma levels of indomethacin are likely to be increased. Therefore, a lower total daily dosage of INDOCIN may produce a satisfactory therapeutic effect. When increases in the dose of INDOCIN are made, they should be made carefully and in small increments.

### **Drug/Laboratory Test Interactions**

False-negative results in the dexamethasone suppression test (DST) in patients being treated with INDOCIN have been reported. Thus, results of the DST should be interpreted with caution in these patients.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

In an 81-week chronic oral toxicity study in the rat at doses up to 1 mg/kg/day, indomethacin had no tumorigenic effect.

Indomethacin produced no neoplastic or hyperplastic changes related to treatment in carcinogenic studies in the rat (dosing period 73-110 weeks) and the mouse (dosing period 62-88 weeks) at doses up to 1.5 mg/kg/day.

Indomethacin did not have any mutagenic effect in *in vitro* bacterial tests (Ames test and *E. coli* with or without metabolic activation) and a series of *in vivo* tests including the host-mediated assay, sex-linked recessive lethals in *Drosophila*, and the micronucleus test in mice.

Indomethacin at dosage levels up to 0.5 mg/kg/day had no effect on fertility in mice in a two generation reproduction study or a two litter reproduction study in rats.



**Pregnancy****Teratogenic Effects. Pregnancy Category C.**

Teratogenic studies were conducted in mice and rats at dosages of 0.5, 1.0, 2.0, and 4.0 mg/kg/day. Except for retarded fetal ossification at 4 mg/kg/day considered secondary to the decreased average fetal weights, no increase in fetal malformations was observed as compared with control groups. Other studies in mice reported in the literature using higher doses (5 to 15 mg/kg/day) have described maternal toxicity and death, increased fetal resorptions, and fetal malformations. Comparable studies in rodents using high doses of aspirin have shown similar maternal and fetal effects. However, animal reproduction studies are not always predictive of human response. There are no adequate and well-controlled studies in pregnant women.

INDOCIN should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

**Nonteratogenic Effects**

Because of the known effects of nonsteroidal anti-inflammatory drugs on the fetal cardiovascular system (closure of ductus arteriosus), use during pregnancy (particularly late pregnancy) should be avoided.

The known effects of indomethacin and other drugs of this class on the human fetus during the third trimester of pregnancy include: constriction of the ductus arteriosus prenatally, tricuspid incompetence, and pulmonary hypertension; non-closure of the ductus arteriosus postnatally which may be resistant to medical management; myocardial degenerative changes, platelet dysfunction with resultant bleeding, intracranial bleeding, renal dysfunction or failure, renal injury/dysgenesis which may result in prolonged or permanent renal failure, oligohydramnios, gastrointestinal bleeding or perforation, and increased risk of necrotizing enterocolitis.

In rats and mice, 4.0 mg/kg/day given during the last three days of gestation caused a decrease in maternal weight gain and some maternal and fetal deaths. An increased incidence of neuronal necrosis in the diencephalon in the live-born fetuses was observed. At 2.0 mg/kg/day, no increase in neuronal necrosis was observed as compared to the control groups. Administration of 0.5 or 4.0 mg/kg/day during the first three days of life did not cause an increase in neuronal necrosis at either dose level.

**Labor and Delivery**

In rat studies with NSAIDs, as with other drugs known to inhibit prostaglandin synthesis, an increased incidence of dystocia, delayed parturition, and decreased pup survival occurred. The effects of INDOCIN on labor and delivery in pregnant women are unknown.

**Use in Nursing Mothers**

INDOCIN is excreted in the milk of lactating mothers. INDOCIN is not recommended for use in nursing mothers.

**Pediatric Use**

Safety and effectiveness in pediatric patients 14 years of age and younger has not been established.

INDOCIN should not be prescribed for pediatric patients 14 years of age and younger unless toxicity or lack of efficacy associated with other drugs warrants the risk.

In experience with more than 900 pediatric patients reported in the literature or to the manufacturer who were treated with Capsules INDOCIN, side effects in pediatric patients were comparable to those reported in adults. Experience in pediatric patients has been confined to the use of Capsules INDOCIN.

If a decision is made to use indomethacin for pediatric patients two years of age or older, such patients should be monitored closely and periodic assessment of liver function is recommended. There have been cases of hepatotoxicity reported in pediatric patients with juvenile rheumatoid arthritis, including fatalities. If indomethacin treatment is instituted, a suggested starting dose is 1-2 mg/kg/day given in divided doses. Maximum daily dosage should not exceed 3 mg/kg/day or 150-200 mg/day, whichever is less. Limited data are available to support the use of a maximum daily dosage of 4 mg/kg/day or 150-200 mg/day, whichever is less. As symptoms subside, the total daily dosage should be reduced to the lowest level required to control symptoms, or the drug should be discontinued.

**Geriatric Use**

As with any NSAID, caution should be exercised in treating the elderly (65 years and older) since advancing age appears to increase the possibility of adverse reactions (see **WARNINGS**,

**Gastrointestinal Effects Risk of Ulceration, Bleeding, and Perforation** and **DOSAGE AND**

**ADMINISTRATION).** Elderly patients seem to tolerate ulceration or bleeding less well than other individuals and many spontaneous reports of fatal GI events are in this population (see **WARNINGS, Gastrointestinal Effects - Risk of Ulceration, Bleeding, and Perforation**).

Indomethacin may cause confusion or, rarely, psychosis (see **ADVERSE REACTIONS**); physicians should remain alert to the possibility of such adverse effects in the elderly.

This drug is known to be substantially excreted by the kidney and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection and it may be useful to monitor renal function (see **WARNINGS, Renal Effects**).

## ADVERSE REACTIONS

In a gastroscopic study in 45 healthy subjects, the number of gastric mucosal abnormalities was significantly higher in the group receiving Capsules INDOCIN than in the group taking Suppositories INDOCIN or placebo.

In a double-blind comparative clinical study involving 175 patients with rheumatoid arthritis, however, the incidence of upper gastrointestinal adverse effects with Suppositories or Capsules INDOCIN was comparable. The incidence of lower gastrointestinal adverse effects was greater in the suppository group.

The adverse reactions for Capsules INDOCIN listed in the following table have been arranged into two groups: (1) incidence greater than 1%; and (2) incidence less than 1%. The incidence for group (1) was obtained from 33 double-blind controlled clinical trials reported in the literature (1,092 patients). The incidence for group (2) was based on reports in clinical trials, in the literature, and on voluntary reports since marketing. The probability of a causal relationship exists between INDOCIN and these adverse reactions, some of which have been reported only rarely.

The adverse reactions reported with Capsules INDOCIN may occur with use of the suppositories. In addition, rectal irritation and tenesmus have been reported in patients who have received the suppositories.

The adverse reactions reported with Capsules INDOCIN may also occur with use of the suspension.

<b>Incidence greater than 1%</b>	<b>Incidence less than 1%</b>	
<b>GASTROINTESTINAL</b>		
nausea** with or without vomiting	anorexia	gastrointestinal bleeding without obvious ulcer formation and perforation of pre-existing sigmoid lesions (diverticulum, carcinoma, etc.)
dyspepsia** (including indigestion, heartburn and epigastric pain)	bloating (includes distension)	development of ulcerative colitis and regional ileitis
diarrhea	flatulence	ulcerative stomatitis
abdominal distress or pain	peptic ulcer	toxic hepatitis and jaundice (some fatal cases have been reported)
constipation	gastroenteritis	intestinal strictures (diaphragms)
	rectal bleeding	
	proctitis	
	single or multiple ulcerations, including perforation and hemorrhage of the esophagus, stomach, duodenum or small and large intestines	
	intestinal ulceration associated with stenosis and obstruction	

INDOCIN<sup>→</sup> (Indomethacin)

9676003

*CENTRAL NERVOUS SYSTEM*

headache (11.7%)  
dizziness\*\*  
vertigo  
somnolence  
depression and fatigue  
(including malaise and  
listlessness)

anxiety (includes  
nervousness)  
muscle weakness  
involuntary muscle  
movements  
insomnia  
muzziness  
psychic disturbances  
including psychotic  
episodes  
mental confusion  
drowsiness

light-headedness  
syncope  
paresthesia  
aggravation of epilepsy  
and parkinsonism  
depersonalization  
coma  
peripheral neuropathy  
convulsion  
dysarthria

*SPECIAL SENSES*

tinnitus

ocular — corneal  
deposits and retinal  
disturbances,  
including those of the  
macula, have been  
reported in some  
patients on prolonged  
therapy with INDOCIN

blurred vision  
diplopia  
hearing disturbances,  
deafness

*CARDIOVASCULAR*

none

hypertension  
hypotension  
tachycardia  
chest pain

congestive heart failure  
arrhythmia; palpitations

*METABOLIC*

none edema

weight gain  
fluid retention  
flushing or sweating

hyperglycemia  
glycosuria  
hyperkalemia

*INTEGUMENTARY*

none pruritus

rash; urticaria  
petechiae or ecchymosis

exfoliative dermatitis  
erythema nodosum  
loss of hair  
Stevens-Johnson  
syndrome  
erythema multiforme  
toxic epidermal necrolysis

*HEMATOLOGIC*

none leukopenia

bone marrow depression  
anemia secondary to  
obvious or occult  
gastrointestinal  
bleedingaplastic anemia  
hemolytic anemia  
agranulocytosis  
thrombocytopenic purpura  
disseminated intravascular  
coagulation**HYPERSENSITIVITY**

none

acute anaphylaxis  
acute respiratory distress  
rapid fall in blood  
pressure resembling a  
shock-like state  
angioedemadyspnea  
asthma  
purpura  
angiitis  
pulmonary edema  
fever**GENITOURINARY**

none hematuria

vaginal bleeding  
proteinuria  
nephrotic syndrome  
interstitial nephritisBUN elevation  
renal insufficiency,  
including renal failure**MISCELLANEOUS**

none epistaxis

breast changes,  
including enlargement  
and tenderness, or  
gynecomastia

---

\*\* Reactions occurring in 3% to 9% of patients treated with INDOCIN. (Those reactions occurring in less than 3% of the patients are unmarked.)

---

**Causal relationship unknown:** Other reactions have been reported but occurred under circumstances where a causal relationship could not be established. However, in these rarely reported events, the possibility cannot be excluded. Therefore, these observations are being listed to serve as alerting information to physicians:

*Cardiovascular:* Thrombophlebitis

*Hematologic:* Although there have been several reports of leukemia, the supporting information is weak

*Genitourinary:* Urinary frequency.

A rare occurrence of fulminant necrotizing fasciitis, particularly in association with Group A  $\beta$  hemolytic streptococcus, has been described in persons treated with non-steroidal anti-inflammatory agents, including indomethacin, sometimes with fatal outcome (see also **PRECAUTIONS, General**).

**OVERDOSAGE**

The following symptoms may be observed following overdosage: nausea, vomiting, intense headache, dizziness, mental confusion, disorientation, or lethargy. There have been reports of paresthesias, numbness, and convulsions.

Treatment is symptomatic and supportive. The stomach should be emptied as quickly as possible if the ingestion is recent. If vomiting has not occurred spontaneously, the patient should be induced to vomit with syrup of ipecac. If the patient is unable to vomit, gastric lavage should be performed. Once the stomach has been emptied, 25 or 50 g of activated charcoal may be given. Depending on the condition of

the patient, close medical observation and nursing care may be required. The patient should be followed for several days because gastrointestinal ulceration and hemorrhage have been reported as adverse reactions of indomethacin. Use of antacids may be helpful.

The oral LD<sub>50</sub> of indomethacin in mice and rats (based on 14 day mortality response) was 50 and 12 mg/kg, respectively.

## DOSAGE AND ADMINISTRATION

Carefully consider the potential benefits and risks of INDOCIN and other treatment options before deciding to use INDOCIN. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

After observing the response to initial therapy with INDOCIN, the dose and frequency should be adjusted to suit an individual patient's needs.

INDOCIN is available as 25 and 50 mg Capsules INDOCIN, Oral Suspension INDOCIN, containing 25 mg of indomethacin per 5 mL, and 50 mg Suppositories INDOCIN for rectal use.

Adverse reactions appear to correlate with the size of the dose of INDOCIN in most patients but not all. Therefore, every effort should be made to determine the smallest effective dosage for the individual patient.

### *Pediatric Use*

INDOCIN ordinarily should not be prescribed for pediatric patients 14 years of age and under (see

## **PRECAUTIONS, Pediatric Use**).

### *Adult Use*

Dosage Recommendations for Active Stages of the Following:

1. Moderate to severe rheumatoid arthritis including acute flares of chronic disease; moderate to severe ankylosing spondylitis; and moderate to severe osteoarthritis.

Suggested Dosage:

Capsules INDOCIN 25 mg b.i.d. or t.i.d. If this is well tolerated, increase the daily dosage by 25 or by 50 mg, if required by continuing symptoms, at weekly intervals until a satisfactory response is obtained or until a total daily dose of 150-200 mg is reached. **DOSES ABOVE THIS AMOUNT GENERALLY DO NOT INCREASE THE EFFECTIVENESS OF THE DRUG.**

In patients who have persistent night pain and/or morning stiffness, the giving of a large portion, up to a maximum of 100 mg, of the total daily dose at bedtime, either orally or by rectal suppositories, may be helpful in affording relief. The total daily dose should not exceed 200 mg. In acute flares of chronic rheumatoid arthritis, it may be necessary to increase the dosage by 25 mg or, if required, by 50 mg daily.

If minor adverse effects develop as the dosage is increased, reduce the dosage rapidly to a tolerated dose and **OBSERVE THE PATIENT CLOSELY.**

If severe adverse reactions occur, **STOP THE DRUG.** After the acute phase of the disease is under control, an attempt to reduce the daily dose should be made repeatedly until the patient is receiving the smallest effective dose or the drug is discontinued.

Careful instructions to, and observations of, the individual patient are essential to the prevention of serious, irreversible, including fatal, adverse reactions.

As advancing years appear to increase the possibility of adverse reactions, INDOCIN should be used with greater care in the elderly (see **PRECAUTIONS, Geriatric Use**).

2. Acute painful shoulder (bursitis and/or tendinitis).

Initial Dose:

75-150 mg daily in 3 or 4 divided doses.

The drug should be discontinued after the signs and symptoms of inflammation have been controlled for several days. The usual course of therapy is 7-14 days.

3. Acute gouty arthritis.

Suggested Dosage:

Capsules INDOCIN 50 mg t.i.d. until pain is tolerable. The dose should then be rapidly reduced to complete cessation of the drug. Definite relief of pain has been reported within 2 to 4 hours.

Tenderness and heat usually subside in 24 to 36 hours, and swelling gradually disappears in 3 to 5 days.

**HOW SUPPLIED**

No. 3316 — Capsules INDOCIN, 25 mg are opaque blue and white capsules, coded INDOCIN and MSD 25. They are supplied as follows:

**NDC** 0006-0025-68 bottles of 100

**NDC** 0006-0025-82 bottles of 1000.

No. 3317 — Capsules INDOCIN, 50 mg are opaque blue and white capsules, coded INDOCIN and MSD 50. They are supplied as follows:

**NDC** 0006-0050-68 bottles of 100.

No. 3376 — Oral Suspension INDOCIN, 25 mg per 5 mL, is an off-white suspension with a pineapple coconut mint flavor. It is supplied as follows:

**NDC** 0006-3376-66 in bottles of 237 mL.

No. 3354 — Suppositories INDOCIN, 50 mg each, are white, opaque, rectal suppositories and are supplied as follows:

**NDC** 0006-0150-30, boxes of 30.


**Storage**

Store Oral Suspension INDOCIN below 30°C (86°F). Avoid temperatures above 50°C (122°F). Protect from freezing.

Store Suppositories INDOCIN below 30°C (86°F). Avoid transient temperatures above 40°C (104°F).


---

Suppositories INDOCIN<sup>®</sup> are distributed by:

 **MERCK & CO., INC.**, Whitehouse Station, NJ 08889, USA

Manufactured by:  
MERCK SHARP & DOHME  
(Italia) S.p.A.  
27100 — Pavia, Italy

Capsules and Oral Suspension INDOCIN<sup>®</sup> are distributed and manufactured by:

 **MERCK & CO., INC.**, Whitehouse Station, NJ 08889, USA

Issued March 2007  
Printed in USA